

# On the Independence Numbers of a Matroid

YAHYA OULD HAMIDOUNE AND ISABELLE SALAÛN

*Université Pierre et Marie Curie, UER 48, ER Combinatoire,  
4 Place Jussieu, 75230 Paris, France*

*Communicated by W. R. Pulleyblank*

Received November 23, 1987

Given a finite subset  $E$  of a vector space of dimension 4, the number of  $k$ -independent subsets of  $E$  will be denoted by  $I_k$ . We prove that  $kI_k^2 \geq (k+1)I_{k-1}I_{k+1} + I_{k-1}I_k$ . The equality holds if and only if all 4-subsets of  $E$  are independent. We prove this relation for matroids of rank 4. In particular we prove Mason's conjecture on the independence numbers of a matroid for  $k=3$ .

© 1989 Academic Press, Inc.

## 1. INTRODUCTION

We use the notations and terminology of Welsh [7].

Let  $M$  be a matroid. The set of independent  $k$ -sets will be denoted by  $\mathcal{I}_k$  and  $|\mathcal{I}_k|$  by  $I_k$ .

The following conjectures are well known.

*Conjecture 1.1* (Welsh [7]). The sequence  $\{I_k\}$  is unimodal.

*Conjecture 1.2* (Mason [3]).  $I_k^2 \geq I_{k+1}I_{k-1}$ .

*Conjecture 1.3* (Mason [3]).  $k(n-k)I_k^2 \geq (k+1)(n-k+1)I_{k+1}I_{k-1}$ , where  $n$  is the number of elements of  $M$ .

Conjecture 1.2 is implied by Dowling's partition conjecture [1] and by a property suggested by Stanley [6]. It is proved for  $k \leq 7$  by Dowling [1]. Seymour proved Conjecture 1.3 for matroids without a circuit  $C$  such that  $3 \leq |C| \leq k-1$  [4].

We propose the following conjecture.

*Conjecture 1.4.*

$$kI_k^2 \geq (k+1)I_{k-1}I_{k+1} + I_{k-1}I_k.$$

Conjecture 1.4 implies Conjecture 1.3 with a strict inequality unless every  $(k+1)$ -subset is independent. This follows immediately from the obvious inequality  $(k+1)I_{k+1} \leq (n-k)I_k$  which is strict unless every  $(k+1)$ -subset is independent.

## 2. NOTATIONS AND PRELIMINAIRES

Let  $G$  be a graph and let  $x$  be a vertex of  $G$ . The set of vertices adjacent to  $x$  will be denoted by  $\Gamma(x)$ . The degree of  $x$  will be denoted by  $d(x)$ . All graphs considered are assumed to be without loops or multiple edges.

Let  $G = (V, W; E)$  be a bipartite graph without isolated vertices and let  $x \in V \cup W$ . We define

$$\theta_G(x) = \sum_{y \in \Gamma(x)} (d(y))^{-1}.$$

LEMMA 2.1 (Seymour [5]).

$$\sum_{x \in V} \theta(x) = |W|.$$

*Proof.*

$$\sum_{x \in V} \theta(x) = \sum_{x \in V} \sum_{y \in \Gamma(x)} (d(y))^{-1} = \sum_{y \in W} \sum_{x \in \Gamma(y)} (d(y))^{-1} = |W|. \quad \blacksquare$$

This lemma is stated by Seymour [5] in a particular case.

Let  $M$  be a matroid and let

$$E = \{ \{X, Y\} \mid X \in \mathcal{J}_{k-1}, Y \in \mathcal{J}_k \text{ and } X \subset Y \}.$$

The bipartite graph  $(\mathcal{J}_{k-1}, \mathcal{J}_k; E)$  will be denoted by  $G_k(M)$ . We will write  $\Gamma_k$  (resp.  $\theta_k$ ) for  $\Gamma_{G_k}$  (resp.  $\theta_{G_k}$ ). The reference to  $M$  will be implicit in general. For  $A \subset E(M)$ , the matroid obtained by restriction to  $A$  (resp. contracting  $A$ ) will be denoted by  $M[A]$  (resp.  $M/A$ ).

We see easily that  $\theta_k(X) = d(X)/k$ , for every  $X \in \mathcal{J}_{k-1}$ .

LEMMA 2.2 (Seymour [5]). *Let  $G = (V, W; E)$  be a bipartite graph such that for every  $Y \subset W$   $|\Gamma(Y)| |W| \geq |Y| |V|$ . Then there is a mapping  $f: V \times W \rightarrow \mathbb{N}$  such that  $f(x, y) = 0$  for any two non-adjacent vertices  $x$  and  $y$  and for each  $y \in W$ ,  $\sum_{x \in V} f(x, y) = |V|$  and for each  $x \in V$ ,  $\sum_{y \in W} f(x, y) = |W|$ .*

This lemma is an easy generalization of the exercise in Seymour [5, p. 20, (2.4)].

LEMMA 2.3. *Let  $M$  be a matroid and let  $k$  be an integer,  $1 \leq k \leq r(M) - 1$ . Let  $(X, Y) \in \mathcal{I}_{k-1} \times \mathcal{I}_k$  be an edge of  $G_k$ . Then  $k\theta_k(X) \geq (k+1)\theta_{k+1}(Y) + 1$ . Moreover the inequality is strict for at least one pair unless the matroid contains no circuit of order less than  $k+2$ .*

*Proof.* Let  $\{b\} = Y - X$ . For every  $Z \in \Gamma_{k+1}(Y)$ , let  $f(Z) = Z - \{b\}$ . The mapping  $f$  is clearly an injection from  $\Gamma_{k+1}(Y)$  to  $\Gamma_k(X) - \{Y\}$ . Therefore  $d_{k+1}(Y) \leq d_k(X) - 1$ .

Since  $\theta_k(X) = d(X)/k$  and  $\theta_{k+1}(Y) = d(Y)/(k+1)$  we have  $k\theta_k(X) \geq (k+1)\theta_{k+1}(Y) + 1$ .

Suppose that there is a circuit  $C$  of  $M$  with cardinality less than  $k+2$ . Let  $B$  be a basis such that  $|B \cap C| = |C| - 1$ . Let  $Y$  be a  $k$ -subset of  $B$  such that  $|Y \cap C| = |C| - 1$  and  $X = Y - b$ , where  $b \in C \cap Y$ . Let  $c \in C - Y$ . It is clear that  $X \cup \{c\} \in \Gamma_k(X) - f(\Gamma_{k+1}(Y))$ . ■

### 3. A STRONG PROPERTY

Let  $M$  be a matroid such that  $E(M) = A \cup B$ , where  $A \in \mathcal{I}_{k-1}$  and  $B \in \mathcal{I}_k$  and  $A \cap B = \emptyset$ . The complement of  $X$  will be denoted by  $\bar{X}$ .

We take  $\mathcal{I}_{k-1} = \{X \in \mathcal{I}_{k-1} \mid \bar{X} \in \mathcal{I}_k\}$  and  $\mathcal{I}_k = \{X \in \mathcal{I}_k \mid \bar{X} \in \mathcal{I}_{k-1}\}$ .

The subgraph of  $G_k(M)$  induced on  $\mathcal{I}_{k-1} \cup \mathcal{I}_k$  will be denoted by  $H_k(M)$ . It is clear that  $|\mathcal{I}_{k-1}| = |\mathcal{I}_k|$ . We define  $\pi_{A,B} = \{(S, T) \in \mathcal{I}_{k-1} \times \mathcal{I}_k \mid S \cup T = A \cup B \text{ and } S \cap T = A \cap B\}$ .

It is clear that  $\{\pi_{X,Y} \mid (X, Y) \in \mathcal{I}_{k-1} \times \mathcal{I}_k\}$  is a partition of  $\mathcal{I}_{k-1} \times \mathcal{I}_k$ .

LEMMA 3.1. *Let  $(A, B) \in \mathcal{I}_{k-1} \times \mathcal{I}_k$ , and let  $\mathcal{P}$  and  $\mathcal{X}$  be two subsets of  $\mathcal{I}_{k-1}$  and  $\mathcal{I}_k$ , respectively. Let  $F = A \cup B$ . Then  $\pi_{A,B} \cap (\mathcal{P} \times \mathcal{X}) = \pi_{A,B} \cap (\mathcal{P}_F \times \mathcal{X}_F)$ , where  $\mathcal{P}_F = \mathcal{P} \cap \mathcal{I}_{k-1}(M[F])$  and  $\mathcal{X}_F = \mathcal{X} \cap \mathcal{I}_k(M[F])$ .*

This lemma is obvious.

We introduce some notations. Let  $\mathcal{X}$  be a set of subsets of  $E$  and  $C \subset E$ ,

$$\mathcal{X}^C = \{X \in \mathcal{X} \mid C \subset X\}$$

$$\mathcal{X} - C = \{X - C \mid X \in \mathcal{X}\}$$

$$\mathcal{X} + C = \{X \cup C \mid X \in \mathcal{X}\}.$$

LEMMA 3.2. *Let  $(A, B) \in \mathcal{I}_{k-1} \times \mathcal{I}_k$  and let  $\mathcal{X}$  be a subset of  $\mathcal{I}_k$  and  $C = A \cap B$ . Let  $M' = M/C$ . Then  $\Gamma_{k'}(\mathcal{X}^C - C) + C \subset \Gamma_k(\mathcal{X})$ , where  $k' = k - |C|$ .*

This lemma is obvious.

**PROPOSITION 3.3.** *Let  $M$  be a matroid such that  $H_j(N)$  has a perfect matching for every  $j \leq k$  and every minor  $N$  of  $M$  with order  $2j - 1$ . Then for any  $\mathcal{X} \subset \mathcal{I}_k$ ,  $I_k |\Gamma_k(\mathcal{X})| \geq I_{k-1} |\mathcal{X}|$ .*

*Proof.* We shall prove the stronger property

$$|\pi_{A,B} \cap (\mathcal{I}_{k-1} \times \mathcal{X})| \leq |\pi_{A,B} \cap (\Gamma_k(\mathcal{X}) \times \mathcal{I}_k)|, \quad \text{for all } (A, B) \in \mathcal{I}_{k-1} \times \mathcal{I}_k.$$

Suppose the contrary. We may assume that  $k$  is minimal such that there is a matroid  $M$  verifying the conditions of the proposition and  $(A, B) \in \mathcal{I}_{k-1} \times \mathcal{I}_k$  such that  $|\pi_{A,B} \cap (\mathcal{I}_{k-1} \times \mathcal{X})| > |\pi_{A,B} \cap (\Gamma_k(\mathcal{X}) \times \mathcal{I}_k)|$ . We shall also choose  $|E(M)|$  minimal. Take  $F = A \cup B$ ,  $C = A \cap B$ , and  $M' = M/C$ . We prove the following.

(1)  $E(M) = F$ . Lemma 3.1 shows that  $M[F]$  is also a counterexample. Therefore  $E(M) = A \cup B$ , by the minimality of the order of  $M$ .

(2)  $C = \emptyset$ . Suppose the contrary. Let  $A' = A - C$ ,  $B' = B - C$ ,  $k' = k - |C|$ , and  $\mathcal{X}' = \mathcal{X}^C - C$ . Since  $H_{k'}(M')$  has a matching and  $k' < k$

$$|\pi_{A',B'} \cap (\mathcal{I}'_{k'-1} \times \mathcal{X}')| \leq |\pi_{A',B'} \cap (\Gamma'_{k'}(\mathcal{X}') \times \mathcal{I}'_{k'})|.$$

It follows that

$$|\pi_{A',B'} \cap (\mathcal{I}'_{k'-1} \times \mathcal{X}') + (C, C)| \leq |\pi_{A',B'} \cap (\Gamma'_{k'}(\mathcal{X}') \times \mathcal{I}'_{k'}) + (C, C)|.$$

Therefore using Lemma 3.2,

$$|\pi_{A,B} \cap ((\mathcal{I}'_{k'-1} + C) \times \mathcal{X}^C)| \leq |\pi_{A,B} \cap (\Gamma_k(\mathcal{X}) \times \mathcal{I}_k)|.$$

But  $\pi_{A,B} \cap ((\mathcal{I}'_{k'-1} + C) \times \mathcal{X}^C) = \pi_{A,B} \cap (\mathcal{I}_{k-1}(M) \times \mathcal{X})$ . Let  $(X, Y) \in \pi_{A,B} \cap (\mathcal{I}_{k-1}(M) \times \mathcal{X})$ . Then  $X \cap Y = C$ . Hence  $Y \in \mathcal{X}^C$ . It follows that  $(X - C, Y - C) \in \mathcal{I}'_{k'-1}(M') \times \mathcal{X}'$

$$|\pi_{A,B} \cap (\mathcal{I}_{k-1}(M) \times \mathcal{X})| \leq |\pi_{A,B} \cap (\Gamma_k(\mathcal{X}) \times \mathcal{I}_k)|, \quad \text{a contradiction.}$$

(3) Now  $|\pi_{A,B} \cap (\mathcal{I}_{k-1} \times \mathcal{X})| = |\mathcal{X} \cap \mathcal{I}_k|$  and

$$|\pi_{A,B} \cap (\Gamma_k(\mathcal{X}) \times \mathcal{I}_k)| = |\Gamma_k(\mathcal{X}) \cap \mathcal{I}_{k-1}|.$$

But it is easy to check that  $\Gamma_k(\mathcal{X}) \cap \mathcal{I}_{k-1} = \Gamma_k(\mathcal{X} \cap \mathcal{I}_k) \cap \mathcal{I}_{k-1}$ . The existence of a matching in  $H_k(M)$  shows that  $|\Gamma_k(\mathcal{X} \cap \mathcal{I}_k) \cap \mathcal{I}_{k-1}| \geq |\mathcal{X} \cap \mathcal{I}_k|$ . It follows that

$$|\pi_{A,B} \cap (\mathcal{I}_{k-1} \times \mathcal{X})| \leq |\pi_{A,B} \cap (\Gamma_k(\mathcal{X}) \times \mathcal{I}_k)|, \quad \text{a contradiction.} \quad \blacksquare$$

There are graphic matroids  $M$  such that  $H_k(M)$  has no matching. The following example due to M. Las Vergnas improves our previous one.

EXAMPLE (Las Vergnas). Let  $G$  be a graph with edge set  $\{12, 13, 14, 15, 23, 24, 25\}$ .  $H_4(\mathbb{C}(G))$  has no perfect matching.

PROPOSITION 3.4. *Let  $M$  be a matroid such that  $H_j(N)$  has a perfect matching for every  $j \leq k$  and every minor  $N$  of  $M$  with order  $2j-1$ . Then  $kI_k^2 \geq I_{k-1}I_k + (k+1)I_{k-1}I_{k+1}$ . Moreover the inequality is strict unless every  $(k+1)$ -set is independent.*

*Proof.* By Proposition 3.3, for any  $\mathcal{X} \subset \mathcal{I}_k$ ,  $I_k |\Gamma_k(\mathcal{X})| \geq I_{k-1} |\mathcal{X}|$ . By Lemma 2.2, there is a mapping  $f: \mathcal{I}_{k-1} \times \mathcal{I}_k \rightarrow \mathbb{N}$  such that  $f(X, Y) = 0$  if  $X$  is not contained in  $Y$ ,  $\sum_X f(X, Y) = I_{k-1}$ , and  $\sum_Y f(X, Y) = I_k$ . Using Lemma 2.3, we obtain

$$kf(X, Y)\theta_k(X) \geq (k+1)f(X, Y)\theta_{k+1}(Y) + f(X, Y) \quad (1)$$

Therefore

$$k \sum_X \left( \sum_Y f(X, Y) \right) \theta_k(X) \geq (k+1) \sum_Y \left( \sum_X f(X, Y) \right) \theta_{k+1}(Y) + \sum_X \sum_Y f(X, Y).$$

By Lemma 2.1 and Lemma 2.2, we obtain

$$kI_k^2 \geq (k+1)I_{k-1}I_{k+1} + I_{k-1}I_k.$$

If  $M$  contains a circuit of order  $\leq k+1$ , Lemma 2.3 shows that (1) is strict for a least one pair. ■

#### 4. A SPECIAL CASE

LEMMA 4.1. *Let  $M$  be a matroid such that  $E(M)$  is the disjoint union of a  $k$ -independent set and a  $(k-1)$ -independent set. Then  $H_k(M)$  has a perfect matching if and only if for every  $\mathcal{X} \subset \mathcal{I}_{k-1}$*

$$|\{Y \in \Gamma_k(\mathcal{X}) \mid \bar{Y} \in \mathcal{I}_{k-1} - \mathcal{X}\}| \geq |\{X \in \mathcal{X} \mid \bar{X} \in \mathcal{I}_k - \Gamma_k(\mathcal{X})\}|.$$

*Proof.* By Hall's condition, there is a perfect matching if and only if for every  $\mathcal{X} \subset \mathcal{I}_{k-1}$ ,

$$\begin{aligned} & |\{Y \in \Gamma_k(\mathcal{X}) \mid \bar{Y} \in \mathcal{I}_{k-1} - \mathcal{X}\}| + |\{Y \in \Gamma_k(\mathcal{X}) \mid \bar{Y} \in \mathcal{X}\}| \\ & \geq |\{X \in \mathcal{X} \mid \bar{X} \in \mathcal{I}_k - \Gamma_k(\mathcal{X})\}| + |\{X \in \mathcal{X} \mid \bar{X} \in \Gamma_k(\mathcal{X})\}|. \end{aligned}$$

But  $|\{Y \in \Gamma_k(\mathcal{X}) \mid \bar{Y} \in \mathcal{X}\}| = |\{X \in \mathcal{X} \mid \bar{X} \in \Gamma_k(\mathcal{X})\}|$ . ■

LEMMA 4.2. *Let  $M$  be a matroid such that  $E(M)$  is the disjoint union of a  $k$ -independent set and a  $(k-1)$ -independent set. Suppose that there is an element  $a$  of  $M$  contained in all  $k$ -independent sets. Then  $H_k(M)$  has a perfect matching.*

*Proof.* Let  $X$  be an element of  $\mathcal{I}_{k-1}(M)$ . Then  $a \in \bar{X}$ . It is easy to see that  $X \cup \{a\}$  is a  $k$ -independent set containing  $X$ . Moreover  $\overline{X \cup \{a\}} = \bar{X} - a$ . Hence  $X \cup \{a\} \in \mathcal{I}_k(M)$ . It follows that  $\{(X, X \cup \{a\}); X \in \mathcal{I}_{k-1}(M)\}$  is a perfect matching. ■

LEMMA 4.3. *Let  $M$  be a matroid such that  $E(M)$  is the disjoint union of a  $k$ -independent set and a  $(k-1)$ -independent set. Suppose that  $2 \leq k \leq 3$ . Then  $H_k(M)$  has a perfect matching.*

*Proof.* If  $k=2$  then the matroid is of order 3. If every 2-set is independent, the perfect matching is obvious. Otherwise there is an element contained in all 2-independent sets. The matching exists by Lemma 4.2.

Suppose the result false for  $k=3$ . The matroid is of order 5. There is no element contained in all 3-independent sets, by Lemma 4.2. Therefore every 2-independent set is contained in at least two distinct 3-independent sets. Take  $\mathcal{X}_1 = \{Y \in \Gamma_k(\mathcal{X}) \mid \bar{Y} \in \mathcal{I}_{k-1} - \mathcal{X}\}$  and  $\mathcal{X}_2 = \{X \in \mathcal{X} \mid \bar{X} \in \mathcal{I}_k - \Gamma_k(\mathcal{X})\}$ . The non-existence of a perfect matching and Lemma 4.1 show that  $|\mathcal{X}_1| < |\mathcal{X}_2|$ . Consider the bipartite graph  $R$  defined on  $\mathcal{X}_1 \cup \mathcal{X}_2$  by inclusion. We have seen that every  $X \in \mathcal{X}_2$  is contained in two 3-independent sets  $X \cup \{a\}$  and  $X \cup \{b\}$ . Suppose that  $X \cup \{a\} \notin \mathcal{X}_1$ . Then  $\bar{X} - a \in \mathcal{X}$ . It follows that  $\bar{X} \in \Gamma_k(\mathcal{X})$ , contradicting the definition of  $\mathcal{X}_2$ . Hence  $d(X) \geq 2$  in  $R$ . An easy argument shows that  $\mathcal{X}_1$  contains at least two vertices of degree 3. The neighbourhood of these sets contains two disjoint sets  $U, V \in \mathcal{X}_2$ . Now  $U \subset \bar{V}$ . Hence  $\bar{V} \in \Gamma_k(\mathcal{X})$ , contradicting the definition of  $\mathcal{X}_2$ . ■

PROPOSITION 4.4. *Let  $M$  be a matroid and  $k$  be an integer  $\leq 3$ . Then  $kI_k^2 \geq I_{k-1}I_k + (k+1)I_{k-1}I_{k+1}$ . Moreover the inequality is strict unless every  $(k+1)$ -set is independent.*

*Proof.* This is an easy application of Proposition 3.4 and Lemma 4.3. ■

COROLLARY 4.5. *Let  $M$  be a matroid of order  $n$ . Then  $I_3^2 \geq (4(n-2)/3(n-3))I_2I_4$ . Moreover the inequality is strict unless the matroid contains no circuit of order  $\leq 4$ .*

The proof results from the remark given after Conjecture 1.4.

## REFERENCES

1. T. A. DOWLING, On the independent set numbers of a finite matroid, *Ann. Dis. Math.* **8** (1980), 21–28.
2. C. MAHONEY, On the unimodality of the independent set numbers of a class of matroids, *J. Combin. Theory Ser. B* **39** (1985), 77–85.
3. J. H. MASON, Matroids: Unimodal conjectures and Motzkin's theorem, in "Combinatorics" (D. J. A. Welsh and D. R. Woodall, Eds.), pp. 207–221, Oxford Univ. Press, Oxford, 1972.
4. P. D. SEYMOUR, "A unimodality problem in matroids," pp. 66–86, Ph.D. thesis, Oxford, 1975.
5. P. D. SEYMOUR, On the points-lines-planes conjecture, *J. Combin. Theory Ser. B* **33** (1982), 17–26.
6. R. P. STANLEY, Two combinatorial applications of the Alexandroff Fenchel inequalities, *J. Combin. Theory Ser. A* **31** (1981), 56–65.
7. D. J. A. WELSH, "Matroid Theory," Academic Press, London/New York, 1976 .